



# The Scattering Amplitude for the Shrödinger Operator in a Layer

Michel Cristofol, Patricia Gaitan

## ► To cite this version:

Michel Cristofol, Patricia Gaitan. The Scattering Amplitude for the Shrödinger Operator in a Layer. 2003. hal-00128231

**HAL Id: hal-00128231**

**<https://hal.science/hal-00128231>**

Preprint submitted on 31 Jan 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Scattering Amplitude for the Schrödinger Operator in a Layer

Michel Cristofol

Patricia Gaitan\*

## Abstract

For a layer with compactly supported inhomogeneity we give an appropriate definition of the scattering amplitude. Then we recover the relation between the far field pattern and the scattering amplitude.

## 1 Introduction

The scattering amplitude is a wellknown way to solve some inverse scattering problem. In the litterature we actually find the definition of the scattering amplitude in the whole homogeneous perturbed space [5],[3] or half space [1]. For a stratified space, this definition is more complicated [2]. The situation in the layer, not yet studied, introduces also difficulties. In this paper we are looking for a definition of the scattering amplitude in the layer. Such wave guides can modelize problems of wave propagation in geophysic, underwater acoustic and so on.

One consider the Schrödinger operator  $A = -\Delta + V(X)$  in  $L^2(\Omega)$  which derives from  $A_0 = -\Delta$  in  $L^2(\Omega)$ , where  $\Omega = \mathbb{R}^n \times (0, \pi)$ ,  $n \geq 1$  and  $D(A) = D(A_0) = H_0^1(\Omega) \cap H^2(\Omega)$ . We study the boundary value problem:

$$\begin{cases} (-\Delta + V(X))u(X) = \lambda u(X) & \text{in } L^2(\Omega), \\ u(X)|_{\partial\Omega} = 0. \end{cases} \quad (1.1)$$

We define the scattering amplitude and the outgoing solutions of (1.1). A work in progress uses the scattering amplitude to establish uniqueness theorem for the potential  $V(X)$ .

The paper is organized as follow. In section 2, we carry out the generalized eigenfunctions of  $A$  and we define the wave operators. We state results on the existence and completeness of these operators. This allows us to prove that the system of generalized eigenfunctions of  $A$  is dense. In section 3, we define the scattering operator, the scattering matrix and finally the scattering amplitude. In section 4, we give an asymptotic development of the generalized eigenfunctions of  $A$  and we link it with the scattering amplitude.

## 2 Spectral Study of $A$

### 2.1 Generalized Eigenfunctions of $A_0$

By a standard Fourier technique, we obtain the generalized eigenfunctions  $\psi_j^0$  of  $A_0$  associated to the eigenvalues  $\lambda_j(|p|)$  in the following form

$$\psi_j^0(x', y, p) = (2\pi)^n \sqrt{\frac{2}{\pi}} e^{ix' \cdot p} \sin(jy), \quad \text{and} \quad \lambda_j(|p|) = j^2 + |p|^2, \quad j \geq 1,$$

with  $X = (x', y)$ ,  $x' = (x_1, \dots, x_n) \in \mathbb{R}^n$ ,  $y \in (0, \pi)$  and  $p \in \mathbb{R}^n$ .

This family of generalized eigenfunctions  $\{\psi_j^0\}_{j \geq 1}$  is dense in  $L^2(\Omega)$ . Then to obtain a spectral representation

---

\*Centre de Mathématiques et d'Informatique, Université de Provence, 39 rue Joliot Curie, 13453 Marseille cedex 13, France, e-mail :cristo@cmi.univ-mrs.fr and gaitan@cmi.univ-mrs.fr

of  $A_0$ , if  $f \in L^2(\Omega)$ , we define the following limit in  $L^2(\Omega)$  and the spaces  $\mathcal{H}_j$ :

$$\begin{aligned}\widetilde{f}_j(p) &= \lim_{M \rightarrow +\infty} \int_{|x'| < M} \overline{\psi_j^0(x', y, p)} f(x', y) dx' dy, \\ \mathcal{H}_j &= \{u \in L^2(\Omega); \exists \widetilde{f}_j \in L^2(\mathbb{R}^n), u(x', y) = \int_{\mathbb{R}^n} \psi_j^0(x', y, p) \widetilde{f}_j(p) dp\}.\end{aligned}\quad (2.1)$$

The unitary map  $\Phi : f \mapsto (\widetilde{f}_1(p), \widetilde{f}_2(p), \dots)$  from  $L^2(\Omega)$  into  $\mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \dots$  is a spectral representation of  $A_0$  in the following sense

$$\Phi(A_0 f) = (\widetilde{A_0 f_1}, \widetilde{A_0 f_2}, \dots) = (\lambda_1(|p|) \widetilde{f}_1(p), \lambda_2(|p|) \widetilde{f}_2(p), \dots).$$

## 2.2 Generalized Eigenfunctions of $A$

We look for the generalized eigenfunctions of  $A$  in the form:

$$\psi_j^\pm(x', y, p) = \psi_j^0(x', y, p) + w_j^\pm(x', y, p), \quad (2.2)$$

We denote

$$N(\lambda) = \#\{j > 1; j^2 \leq \lambda\}. \quad (2.3)$$

The function  $\psi_j^\pm$  satisfies the eigenvalue equation

$$(-\Delta + V(x', y) - \lambda_j(|p|)) \psi_j^\pm(x', y, p) = 0. \quad (2.4)$$

Then, taking into account the equations (2.2), (2.4) and the limiting absorption principle, we obtain

$$w_j^\pm(x', y, p) = -R^\pm(\lambda_j(|p|))(V\psi_j^0)(x', y, p), \quad (2.5)$$

where  $R^\pm(z)$  is the resolvent of  $A$ . If we introduce the resolvent equation in (2.5), we obtain

$$w_j^\pm(x', y, p) = -R_0^\pm(\lambda_j(|p|))(V\psi_j^0)(x', y, p) + R_0^\pm(\lambda_j(|p|))VR^\pm(\lambda_j(|p|))(V\psi_j^0)(x', y, p). \quad (2.6)$$

## 2.3 Wave Operators

We first consider the wave operators

$$W_\pm = s - \lim_{t \rightarrow \pm\infty} e^{itA} e^{-itA_0}, \quad (2.7)$$

and the scattering operator  $S = W_+^* W_-$ . The wave operators actually exist. The proof is an adaptation of the theorem XIII-31 of [6]. Then the limits (2.7) exist, moreover we obtain the completeness of the wave operators  $W_\pm$  from  $\mathcal{H}_{ac}^0(\Omega) = L^2(\Omega)$  (subspace of absolute continuity of  $A_0$ ) into  $\mathcal{H}_{ac}(\Omega)$  (subspace of absolute continuity of  $A$ ), see [4]. To obtain a spectral representation of  $A$ , if  $f \in L^2(\Omega)$ , we define the following limit in  $L^2(\Omega)$ :

$$\widetilde{f}_j^\pm(p) = \lim_{M \rightarrow +\infty} \int_{|x'| < M} \overline{\psi_j^\pm(x', y, p)} f(x', y) dx' dy,$$

and  $\mathcal{H}_j^\pm = W_\pm \mathcal{H}_j$ , where  $\mathcal{H}_j$  is defined by (2.1). The unitary map  $\Phi^\pm : f \mapsto (\widetilde{f}_1^\pm(p), \widetilde{f}_2^\pm(p), \dots)$  from  $L^2(\Omega)$  to  $\mathcal{H}_1^\pm \oplus \mathcal{H}_2^\pm \oplus \dots$  is a spectral representation of  $A|_{\mathcal{H}_{ac}}$ .

### 3 The Scattering Amplitude in a Layer

We have defined in the previous section the scattering operator  $S = W_+^* W_-$ . Using the relations  $\Phi^+ = \Phi W_-^*$  and  $\Phi^- = \Phi W_+^*$ , the scattering operator can be written

$$S = \Phi^* \Phi^- \Phi^{+*} \Phi. \quad (3.1)$$

In a first step we define the scattering matrix as follows

$$\widetilde{S}u_j(p) = (\widetilde{S}(\lambda_j(|p|))u_j)(\omega) \quad \text{with } \omega \in S^{n-1}. \quad (3.2)$$

Note that  $\widetilde{S}(\lambda) \in \mathcal{L}((L^2(S^{n-1}))^{N(\lambda)})$ . We obtain an  $N(\lambda) \times N(\lambda)$ -matrix, where  $N(\lambda)$  was defined by (2.3). So we have to determine  $\widetilde{S}u_j(p)$ . Using (3.1) we obtain  $\Phi S u = \Phi u + (\Phi^- - \Phi^+) \theta^+ u$  where  $\theta^+ = \Phi^{+*} \Phi$ . Then we have

$$\widetilde{S}u_j(p) = \widetilde{u}_j(p) + \lim_{M \rightarrow +\infty} \int_{|x'| < M, y \in (0, \pi)} [\overline{\psi_j^-(x', y, p) - \psi_j^+(x', y, p)}] \theta^+ u(x', y) dx' dy.$$

After several technical transformations, we obtain:

$$\begin{aligned} \widetilde{S}u_j(p) &= \widetilde{u}_j(p) + \sum_{k^2 < j^2 + |p|^2} i\pi \alpha^{n-2} \\ &\left( \int_{\mathbb{R}^n \times (0, \pi)} V(x', y) \psi_j^0(x', y, p) \int_{S^{n-1}} \overline{\psi_k^0(x', y, \alpha\omega')} \widetilde{u}_k(\alpha\omega') d\omega' dx' dy \right. \\ &\left. - \int_{\mathbb{R}^n \times (0, \pi)} V(x', y) \psi_j^0(x', y, p) \int_{S^{n-1}} [\overline{R^+(\lambda_k(\alpha\omega'))} V \psi_k^0(\cdot, \cdot, \alpha\omega')] (x', y, \alpha\omega') \widetilde{u}_k(\alpha\omega') d\omega' dx' dy, \right. \end{aligned} \quad (3.3)$$

with  $\alpha = (\lambda_j(|p|) - k^2)^{1/2}$ . So taking into account (3.2) and (3.3) we obtain explicetely the scattering matrix and the scattering amplitude  $\mathcal{A}(\omega, \omega', \lambda_j(|p|)) = [\mathcal{A}_{j,k}(\omega, \omega', \lambda_j(|p|))]_{1 \leq j, k \leq N(\lambda)}$  is defined as the kernel of this operator

$$((S(\lambda_j(|p|)) - I) \widetilde{u}_j)(\omega) = \int_{S^{n-1}} \sum_{k^2 < j^2 + |p|^2} \mathcal{A}_{j,k}(\omega, \omega', \lambda_j(|p|)) \widetilde{u}_k(\alpha\omega') d\omega'. \quad (3.4)$$

**Proposition 3.1** *The scattering amplitude is a  $N(\lambda) \times N(\lambda)$ -matrix*

$$\begin{aligned} \mathcal{A}(\omega, \omega', \lambda) &= [\mathcal{A}_{j,k}(\omega, \omega', \lambda_j(|p|))]_{1 \leq j, k \leq N(\lambda)} \\ \mathcal{A}_{j,k}(\omega, \omega', \lambda_j(|p|)) &= \mathcal{B}_{j,k}(\omega, \omega', \lambda_j(|p|)) + \mathcal{C}_{j,k}(\omega, \omega', \lambda_j(|p|)), \\ \mathcal{B}_{j,k} &= i\pi \alpha^{n-2} \int_{\mathbb{R}^n \times (0, \pi)} V(x', y) \psi_j^0(x', y, p) \int_{S^{n-1}} \overline{\psi_k^0(x', y, \alpha\omega')} \widetilde{u}_k(\alpha\omega') d\omega' dx' dy \\ \mathcal{C}_{j,k} &= -i\pi \alpha^{n-2} \int_{\mathbb{R}^n \times (0, \pi)} V(x', y) \psi_j^0(x', y, p) \int_{S^{n-1}} [\overline{R^+(\lambda_k(\alpha\omega'))} V \psi_k^0(\cdot, \cdot, \alpha\omega')] (x', y, \alpha\omega') \widetilde{u}_k(\alpha\omega') d\omega' dx' dy. \end{aligned} \quad (3.5)$$

Remark: It should be mentioned the matrix-form of the scattering amplitude which differs radically from the classical one.

### 4 Asymptotic Behaviour of the Generalized Eigenfunctions

The functions  $w_j^\pm$  see (2.2) are defined by

$$\begin{aligned} w_j^\pm(x', y, p) &= -R_0^\pm(\lambda_j(|p|))(V \psi_j^0)(x', y, p) + R_0^\pm(\lambda_j(|p|))V R^\pm(\lambda_j(|p|))(V \psi_j^0)(x', y, p) \\ &= I_{1,j}(x', y, p) + I_{2,j}(x', y, p). \end{aligned}$$

We are looking for the asymptotic behaviour of each term of this sum. Let us consider the  $k^{th}$  component of the first term

$$[I_{1,j}(x', y, p)]_k = \int_{\mathbb{R}^n} \psi_k^0(x', y, p) \frac{1}{|p'|^2 + k^2 - \lambda_j - i0} [V\widetilde{\psi_j^0(\cdot, p)}](p') dp'.$$

Then

$$[I_{1,j}(x', y, p)]_k = (2\pi)^{-3n} \left(\frac{2}{\pi}\right)^{3/2} \sin(ky) \int_{\mathbb{R}^n} \frac{e^{ix' \cdot p'}}{|p'|^2 + k^2 - \lambda_j - i0} f_1(p') dp'$$

$$\text{with } f_1(p') = \int_{\mathbb{R}^n \times (0, \pi)} e^{i\tilde{x}' \cdot (p - p')} \sin(k\tilde{y}) \sin(j\tilde{y}) V(\tilde{x}', \tilde{y}) d\tilde{x}' d\tilde{y}.$$

Remark: To obtain an asymptotic development of  $[I_{1,j}(x', y, p)]_k$ , we use the lemma A1 of [2]. For this we have to replace  $f_1 \in C^\infty(\mathbb{R}^n)$  by a  $C_0^\infty(\mathbb{R}^n)$  function.

Let  $p'_0 = \alpha\omega'$ , ( $p'_0$  verify  $|p'_0|^2 + k^2 = \lambda_j(|p|)$ ), we define the  $C_0^\infty(\mathbb{R}^n)$  function  $\chi$  equal to 1 in a neighborhood of  $p'_0$  and 0 outside. Then

$$[I_{1,j}(x', y, p)]_k = (2\pi)^{-2n} \left(\frac{2}{\pi}\right)^{3/2} \sin(ky) (J_1(x') + K_1(x'))$$

$$J_1(x') = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \frac{e^{ix' \cdot p'} \chi(p')}{|p'|^2 + k^2 - \lambda_j - i0} f_1(p') dp',$$

$$K_1(x') = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \frac{e^{ix' \cdot p'} (1 - \chi(p'))}{|p'|^2 + k^2 - \lambda_j - i0} f_1(p') dp'.$$

Then, we apply the lemma A1 of [2] and we obtain

$$J_1(x') = C_{p'_0} \frac{e^{i|x'| \cdot |p'_0|}}{|x'|^{\frac{n-1}{2}}} (f_1(p'_0) + \mathcal{O}(|x'|^{-1}))$$

where  $C_{p'_0} = (4\pi)^{-1} \alpha^{\frac{n-3}{4}} (2\pi)^{\frac{3-n}{2}} e^{-i\pi \frac{n-3}{4}}$ . Moreover, we prove that  $K_1(x') = \mathcal{O}(|x'|^{-\frac{n+1}{2}})$ . So the asymptotic behaviour of the first term of  $w_j^+$ , denoted  $\overline{[I_{1,j}(x', y, p)]_k}$ , is given by

$$\begin{aligned} \overline{[I_{1,j}(x', y, p)]_k} &= \left(\frac{2}{\pi}\right)^{\frac{1+5n}{2}} \sin(ky) \alpha^{\frac{n-3}{2}} e^{-i\pi \frac{n-3}{4}} \frac{e^{i|x'| \cdot |p'_0|}}{|x'|^{\frac{n-1}{2}}} \\ &\quad \times \int_{\mathbb{R}^n \times (0, \pi)} e^{i\tilde{x}' \cdot (p - p'_0)} \sin(k\tilde{y}) \sin(j\tilde{y}) V(\tilde{x}', \tilde{y}) d\tilde{x}' d\tilde{y}. \end{aligned}$$

In the same way, we prove that the asymptotic behaviour of the second term of  $w_j^+$ , denoted  $\overline{[I_{2,j}(x', y, p)]_k}$ , is given by

$$\begin{aligned} \overline{[I_{2,j}(x', y, p)]_k} &= \frac{(2\pi)^{\frac{1-3n}{2}}}{\pi} \sin(ky) \alpha^{n-3} V(x', y) e^{-i\pi \frac{n-3}{4}} \frac{e^{i|x'| \cdot |p'_0|}}{|x'|^{\frac{n-1}{2}}} \\ &\quad \times \int_{\mathbb{R}^n \times (0, \pi)} e^{-i\tilde{x}' \cdot p'_0} \sin(k\tilde{y}) R(\lambda_j(|p|)) (V(\cdot, \cdot) \psi_j^0(\cdot, \cdot, p))(\tilde{x}', \tilde{y}, p) d\tilde{x}' d\tilde{y}. \end{aligned}$$

In conclusion, we obtain the asymptotic behaviour of the  $k^{th}$  component of  $w_j^+$ :

$$\overline{[w_j^+(x', y, p)]_k} = \overline{[I_{1,j}(x', y, p)]_k} + \overline{[I_{2,j}(x', y, p)]_k}. \quad (4.1)$$

So we have to link the far field pattern with the scattering amplitude. For this, we compare (3.5) with (4.1).

**Proposition 4.1** *If we set*

$$[G(x', y, p)]_k = \sqrt{\frac{2}{\pi}} (2\pi)^{-\frac{n-1}{2}} \sin(ky) \alpha^{-\frac{n+1}{2}} e^{-i\pi \frac{n-1}{2}} \frac{e^{i|x'|\cdot\alpha}}{|x'|^{\frac{n-1}{2}}},$$

with  $\alpha = (\lambda_j(|p|) - k^2)^{1/2}$ , we can write

$$\psi_j^+(x', y, p) = \psi_j^0(x', y, p) + \sum_{k=1}^{N(\lambda)} \mathcal{A}_{j,k}(\omega, \omega', \lambda_j(|p|)) [G(x', y, p)]_k + o(|x'|^{-\frac{n+1}{2}}).$$

This relation allows us to give a physical sense to the scattering amplitude.

Our goal is to prove that exists an injection between the scattering amplitude and the potential  $V(X)$ . For the layer  $(\mathbb{R}^2 \times (0, \pi))$  [3] prove an uniqueness result but using the Dirichlet to Neumann map. In a strip  $(\mathbb{R} \times (0, \pi))$  does not exist any result. We think that the knowledge of the scattering amplitude at fixed energy is not sufficient to determine the potential. We want to prove that a countable infinity of energies will be sufficient.

## References

- [1] G. ESKIN AND J. RALSTON, *Inverse coefficient problems in perturbed half space*, Inverse Problems, 1999 vol 3, p.683-699.
- [2] J.C. GUILLOT AND J. RALSTON, *Inverse scattering at fixed energy for layered media*, J. Math. Pures Appl., 78, 1999, p.27-48 .
- [3] M. IKEHATA , *Reconstruction of a obstacle from the scattering amplitude at fixed energy*, Inverse Problems, vol 14, (1998) p.949-954.
- [4] W.C. LYFORD, *Spectral Analysis of the Laplacian in Domains with Cylinders*, Math. Ann., (1975) p.213-251.
- [5] R.G. NOVIKOV, *On determination of the Fourier transform of a potential from the scattering amplitude*, Inverse Problems, vol 17, (2001) p.1243-1251.
- [6] M. REED AND B. SIMON, *Methods of Modern Mathematical Physics, vol IV, Analysis of Operators* , Academic Press, 1978.